arXiv:1211.3256v1 [math.NT] 14 Nov 2012

ERGODICITY OF THE ACTION OF K^* ON $\mathbb{A}_K$

JEFFREY C. LAGARIAS AND SERGEY NESHVEYEV

ABSTRACT. Connes gave a spectral interpretation of the critical zeros of zeta- and L-functions for a global field K using a space of square integrable functions on the space $\mathbb{A}_K/K^*$ of adele classes. It is known that for $K=\mathbb{Q}$ the space $\mathbb{A}_K/K^*$ cannot be understood classically, or in other words, the action of $\mathbb{Q}^*$ on $\mathbb{A}_\mathbb{Q}$ is ergodic. We prove that the same is true for any global field K, in both the number field and function field cases.

INTRODUCTION

For a global field K, the space of adeles $\mathbb{A}_K$ appeared explicitly (as "valuation vectors") and the adele class space $\mathbb{A}_K/K^*$ appeared implicitly in the work of Tate [19] on the functional equation for Hecke L-functions. More recently $\mathbb{A}_K/K^*$ was used by Connes [4] to construct a Hilbert space that provides a spectral interpretation of the critical zeros of Hecke L-functions. One of the major difficulties in constructing such a Hilbert space, as well as in trying to prove the Riemann hypotheses following Connes' approach, is that the quotient space $\mathbb{A}_K/K^*$ is quite tricky. The precise statement for $K=\mathbb{Q}$ is that the action of $\mathbb{Q}^*$ on $\mathbb{A}_\mathbb{Q}$ is ergodic, so from the point of view of classical measure theory the space $\mathbb{A}_\mathbb{Q}/\mathbb{Q}^*$ consists of just one point. This is a known byproduct of the analysis of the Bost-Connes system [3, 15]. The aim of this paper is to prove that the same ergodicity holds for K^* acting on $\mathbb{A}_K$ for an arbitrary global field K, i.e., for K an algebraic number field or a one-variable algebraic function field over a finite field.

Partial results in this direction have been already known from the analysis of Bost-Connes type systems for global fields [10, 16, 17]. For example, for number fields by [16, Corollary 3.2] the action of the subgroup of totally positive elements in K^* on $\mathbb{R}^*_+\times\mathbb{A}_{K,f}$, given by $x(t,a)=(N((x))t,xa)$, is ergodic. This result could be used to simplify some of our arguments. However, the simplifications would be non-essential, and roughly amount to reducing the arguments below to the case of Hecke characters $(\mathrm{mod}\ 1)$ instead of using all Hecke characters. We have therefore chosen to prove everything what is needed from scratch. As a consequence our arguments can be used to give different proofs of several results in [3, 10, 16, 17], see Sections 5.4 and 5.5.

1. MAIN RESULT

Let K be a global field. Denote by $\mathbb{A}_K$ the adele ring of K and $\mu_\mathbb{A}$ an additive Haar measure on $\mathbb{A}_K$.

Theorem 1.1. *The action of K^* on $(\mathbb{A}_K,\mu_\mathbb{A})$ by multiplication is ergodic; that is, for every K^*-invariant measurable subset of $\mathbb{A}_K$ either the subset itself or its complement has measure zero.*

In order to shorten the notation we will usually omit explicit mentioning of measures. In particular, by ergodicity of an action we will always mean ergodicity with respect to an appropriate Haar measure, or with respect to the push-forward of such a measure, unless said otherwise. The ring $\mathbb{A}_K$ is not compact, and any choice of additive Haar measure $\mu_\mathbb{A}$ for $\mathbb{A}_K$ will assign infinite measure to the full space $\mathbb{A}_K$. The Haar measure is a product-measure, and we assume for convenience that for every non-Archimedean place v of K the Haar measure on K_v is normalized so that the measure of the maximal compact subring $\mathcal{O}_v\subset K_v$ is equal to 1.

Our first goal is to replace $\mathbb{A}_K$ by a more manageable space and the action of K^* by an action of a group of ideals on it.

Date: November 14, 2012, v2.0.

Research supported in part by NSF grant DMS-1101373 and the Norwegian Research Council.

If K is a number field, denote by $\mathcal{O}$ the ring of integers in K. In the case of function fields we fix a place ∞ of K, and then denote by $\mathcal{O}$ the ring of elements in K that have no poles away from ∞, so $\mathcal{O}=\cap_{v\ne\infty}(K\cap\mathcal{O}_v)$.

For a non-zero ideal $\mathfrak{m}$ in $\mathcal{O}$, denote by $I_\mathfrak{m}$ the group of fractional ideals in $\mathcal{O}$ relatively prime to $\mathfrak{m}$. Denote by V_f the set of finite places of K and by $S(\mathfrak{m})\subset V_f$ the set of prime divisors of $\mathfrak{m}$. Put

$$\mathbb{A}_{S(\mathfrak{m})}:={\prod_{v\in V_f\setminus S(\mathfrak{m})}}' K_v,\ \ \hat{\mathcal{O}}_{S(\mathfrak{m})}:=\prod_{v\in V_f\setminus S(\mathfrak{m})}\mathcal{O}_v$$

Writing $\mathfrak{m}$ as $\prod_{v\in S(\mathfrak{m})}\mathfrak{p}_v^{m_v}$, we put

$$W_\mathfrak{m}:=\hat{\mathcal{O}}^*_{S(\mathfrak{m})}\times\prod_{v\in S(\mathfrak{m})}(1+\hat{\mathfrak{p}}_v^{m_v})\subset\hat{\mathcal{O}}^*:=\prod_{v\in V_f}\mathcal{O}_v^*,$$

where $\hat{\mathfrak{p}}_v$ is the maximal ideal in $\mathcal{O}_v$. Identifying $I_\mathfrak{m}$ with $\mathbb{A}^*_{S(\mathfrak{m})}/\hat{\mathcal{O}}^*_{S(\mathfrak{m})}$ we get actions of $I_\mathfrak{m}$ on $\mathbb{A}^*_K/W_\mathfrak{m}$ and $\mathbb{A}_{S(\mathfrak{m})}/\hat{\mathcal{O}}^*_{S(\mathfrak{m})}$ by multiplication. The Haar measure on $\mathbb{A}_{S(\mathfrak{m})}$, normalized so that the measure of $\hat{\mathcal{O}}_{S(\mathfrak{m})}$ is equal to 1, defines a measure on $\mathbb{A}_{S(\mathfrak{m})}/\hat{\mathcal{O}}^*_{S(\mathfrak{m})}$. The quotient $\mathbb{A}^*_K/K^*W_\mathfrak{m}$ of the idele class group is a locally compact group under multiplication, so it also carries a Haar measure.

Theorem 1.2. *The diagonal action of $I_\mathfrak{m}$ on $(\mathbb{A}^*_K/K^*W_\mathfrak{m})\times(\mathbb{A}_{S(\mathfrak{m})}/\hat{\mathcal{O}}^*_{S(\mathfrak{m})})$, equipped with the product measure, is ergodic for every non-zero ideal $\mathfrak{m}\subset\mathcal{O}$.*

In the remaining part of this section we will show that the statements of Theorems 1.1 and 1.2 are equivalent. In the next two sections we then prove Theorem 1.2 for number fields. In Section 2 we give a preliminary result on asymptotic range of measurable cocycles, and in Section 3 we establish Theorem 1.2. In Section 4 we describe modifications to the proof needed to treat the function field case. Finally in Section 5 we give various remarks.

The compact groups $W_\mathfrak{m}$ form a base of neighbourhoods of the identity in $\hat{\mathcal{O}}^*$. Hence by averaging over $W_\mathfrak{m}$ we can approximate any bounded measurable function on $\mathbb{A}_K$ by a $W_\mathfrak{m}$-invariant function for some $\mathfrak{m}$. It follows that the action of K^* on $\mathbb{A}_K$ is ergodic if and only if the action of K^* on $\mathbb{A}_K/W_\mathfrak{m}$ is ergodic for all $\mathfrak{m}$.

Denote by $K^*_{\mathfrak{m},1}\subset K^*$ the subgroup of elements $a\equiv 1 \mod \mathfrak{m}$, and by $\mathcal{P}_{\mathfrak{m},1}\subset I_\mathfrak{m}$ the subgroup of principal fractional ideals having a generator in $K^*_{\mathfrak{m},1}$. Put $U_{\mathfrak{m},1}:=K^*_{\mathfrak{m},1}\cap\mathcal{O}^*$. Consider the subset

$$X_\mathfrak{m}:=\left(K^*_\infty\times\mathbb{A}_{S(\mathfrak{m})}\times\left(\prod_{v\in S(\mathfrak{m})}(1+\hat{\mathfrak{p}}_v^{m_v})\right)\right)/W_\mathfrak{m}\subset\mathbb{A}_K/W_\mathfrak{m},$$

where for number fields $K_\infty=\prod_{v|\infty}K_v$. This set is $K^*_{\mathfrak{m},1}$-invariant, the set $K^*X_\mathfrak{m}\subset\mathbb{A}_K/W_\mathfrak{m}$ is a subset of full measure, and $xX_\mathfrak{m}\cap X_\mathfrak{m}=\emptyset$ for any $x\in K^*\setminus K^*_{\mathfrak{m},1}$. It follows that the action of K^* on $\mathbb{A}_K/W_\mathfrak{m}$ is ergodic if and only if the action of $K^*_{\mathfrak{m},1}$ on

$$X_\mathfrak{m}\cong K^*_\infty\times\mathbb{A}_{S(\mathfrak{m})}/\hat{\mathcal{O}}^*_{S(\mathfrak{m})}$$

is ergodic, or in other words, the action of $\mathcal{P}_{\mathfrak{m},1}\cong K^*_{\mathfrak{m},1}/U_{\mathfrak{m},1}$ on $K^*_\infty/U_{\mathfrak{m},1}\times\mathbb{A}_{S(\mathfrak{m})}/\hat{\mathcal{O}}^*_{S(\mathfrak{m})}$ is ergodic.

Now for each $\mathfrak{m}$, induce the last action of $\mathcal{P}_{\mathfrak{m},1}$ to an action of $I_\mathfrak{m}$. So consider the space

$$Y_\mathfrak{m}:=I_\mathfrak{m}\times_{\mathcal{P}_{\mathfrak{m},1}}(K^*_\infty/U_{\mathfrak{m},1}\times\mathbb{A}_{S(\mathfrak{m})}/\hat{\mathcal{O}}^*_{S(\mathfrak{m})}),$$

with the action of $I_\mathfrak{m}$ defined by its action on $I_\mathfrak{m}$ in the first factor by translations. This action is ergodic if and only if the action of $\mathcal{P}_{\mathfrak{m},1}$ on $K^*_\infty/U_{\mathfrak{m},1}\times\mathbb{A}_{S(\mathfrak{m})}/\hat{\mathcal{O}}^*_{S(\mathfrak{m})}$ is ergodic. The map

$$I_\mathfrak{m}\times K^*_\infty/U_{\mathfrak{m},1}\times\mathbb{A}_{S(\mathfrak{m})}/\hat{\mathcal{O}}^*_{S(\mathfrak{m})}\to I_\mathfrak{m}\times K^*_\infty/U_{\mathfrak{m},1}\times\mathbb{A}_{S(\mathfrak{m})}/\hat{\mathcal{O}}^*_{S(\mathfrak{m})},\ \ (\mathfrak{a},x,b)\mapsto(\mathfrak{a},x,\mathfrak{a}b),$$

defines a measure class preserving homeomorphism of $Y_\mathfrak{m}$ onto

$$Z_\mathfrak{m}:=(I_\mathfrak{m}\times_{\mathcal{P}_{\mathfrak{m},1}}K^*_\infty/U_{\mathfrak{m},1})\times\mathbb{A}_{S(\mathfrak{m})}/\hat{\mathcal{O}}^*_{S(\mathfrak{m})}.$$

Under this homeomorphism the action of $I_\mathfrak{m}$ on $Y_\mathfrak{m}$ becomes the diagonal action on $Z_\mathfrak{m}$, with the action of $I_\mathfrak{m}$ on $I_\mathfrak{m}\times_{\mathcal{P}_{\mathfrak{m},1}}K^*_\infty/U_{\mathfrak{m},1}$ defined by the action on $I_\mathfrak{m}$ in the first factor by translations.

Finally, the group $I_\mathfrak{m}\times_{\mathcal{P}_{\mathfrak{m},1}}K^*_\infty/U_{\mathfrak{m},1}$ is isomorphic to the quotient of $\mathbb{A}^*_{S(\mathfrak{m})}/\hat{\mathcal{O}}^*_{S(\mathfrak{m})}\times K^*_\infty$ by the subgroup $\{(x^{-1},x)\mid x\in K^*_{\mathfrak{m},1}\}$. Applying the automorphism $(x,y)\mapsto(x,y^{-1})$ of $\mathbb{A}^*_{S(\mathfrak{m})}/\hat{\mathcal{O}}^*_{S(\mathfrak{m})}\times K^*_\infty$ we conclude that $I_\mathfrak{m}\times_{\mathcal{P}_{\mathfrak{m},1}}K^*_\infty/U_{\mathfrak{m},1}$ is isomorphic to the quotient of $\mathbb{A}^*_{S(\mathfrak{m})}/\hat{\mathcal{O}}^*_{S(\mathfrak{m})}\times K^*_\infty$ by $K^*_{\mathfrak{m},1}$ under the diagonal embedding. The group $\mathbb{A}^*_{S(\mathfrak{m})}/\hat{\mathcal{O}}^*_{S(\mathfrak{m})}\times K^*_\infty$ can be identified with

$$H_\mathfrak{m}:=\left(K^*_\infty\times\mathbb{A}^*_{S(\mathfrak{m})}\times\left(\prod_{v\in S(\mathfrak{m})}(1+\hat{\mathfrak{p}}_v^{m_v})\right)\right)/W_\mathfrak{m}\subset\mathbb{A}^*_K/W_\mathfrak{m}.$$

We have $K^*H_\mathfrak{m}=\mathbb{A}^*_K/W_\mathfrak{m}$ and $xH_\mathfrak{m}\cap H_\mathfrak{m}=\emptyset$ for any $x\in K^*\setminus K^*_{\mathfrak{m},1}$. It follows that

$$H_\mathfrak{m}/K^*_{\mathfrak{m},1}\cong\mathbb{A}^*_K/K^*W_\mathfrak{m}.$$

To summarize, we have an isomorphism

$$I_\mathfrak{m}\times_{\mathcal{P}_{\mathfrak{m},1}}K^*_\infty/U_{\mathfrak{m},1}\cong\mathbb{A}^*_K/K^*W_\mathfrak{m}$$

that respects the embeddings of $I_\mathfrak{m}\cong\mathbb{A}^*_{S(\mathfrak{m})}/\hat{\mathcal{O}}^*_{S(\mathfrak{m})}$. Therefore the action of K^* on $\mathbb{A}_K/W_\mathfrak{m}$ is ergodic if and only if the action of on $I_\mathfrak{m}$ on $Z_\mathfrak{m}\cong(\mathbb{A}^*_K/K^*W_\mathfrak{m})\times(\mathbb{A}_{S(\mathfrak{m})}/\hat{\mathcal{O}}^*_{S(\mathfrak{m})})$ is ergodic.

This holds for all $\mathfrak{m}$, so the statements of Theorems 1.1 and 1.2 are indeed equivalent.

2. ASYMPTOTIC RANGE OF A COCYCLE

Let (X,μ) be a standard measure space and $\mathcal{R}$ be an ergodic countable equivalence relation on it, as in Feldman and Moore [5]. Assume c is a 1-cocycle on $\mathcal{R}$ with values in an abelian second countable locally compact group Γ, so c is a measurable map $\mathcal{R}\to\Gamma$ such that $c(x,y)c(y,z)=c(x,z)$ for $x\sim_\mathcal{R}y\sim_\mathcal{R}z$. Associate to c an equivalence relation $\mathcal{R}(c)$ on $\Gamma\times X$ defined by

$$(\beta,x)\sim_{\mathcal{R}(c)}(\gamma,y)\ \text{ iff }\ x\sim_\mathcal{R}y\ \text{ and }\ c(x,y)\beta=\gamma.$$

For a measurable subset $A\subset X$ of positive measure denote by c_A the restriction of c to $\mathcal{R}\cap(A\times A)$. The *asymptotic range* $r^*(c)\subset\Gamma$ of c is defined as the intersection of essential ranges of c_A for all measurable subsets $A\subset X$ of positive measure [5, Definition 8.2]. In other words, $\gamma\in\Gamma$ belongs to $r^*(c)$ if and only if for every neighbourhood U of γ and every measurable subset $A\subset X$ of positive measure there exists a measurable subset $B\subset A$ of positive measure and a measurable map $T\colon B\to A$ with graph in $\mathcal{R}$ such that $c(x,Tx)\in U$ for all $x\in B$. We recall the following facts.

Proposition 2.1. *The asymptotic range has the following properties:*

(i) *the set $r^*(c)\subset\Gamma$ is a closed subgroup;*
(ii) *for any measurable subset $A\subset X$ of positive measure we have $r^*(c_A)=r^*(c)$;*
(iii) *the equivalence relation $\mathcal{R}(c)$ on $\Gamma\times X$ is ergodic if and only if $r^*(c)=\Gamma$.*

Proof. Properties (i) and (ii) easily follow from definitions. Property (iii) is nontrivial, it follows from Theorem 8 in [5] (see also Propositions 8.1 and 8.3 there). ☐

Consider now a particular class of ergodic equivalence relations. Take a sequence $\{(X_n,\mu_n)\}_{n=1}^\infty$ of at most countable probability spaces. Put

$$(X,\mu):=\prod_{n=1}^\infty(X_n,\mu_n).$$

Define an equivalence relation $\mathcal{R}$ on X by

$$x\sim_\mathcal{R}y\ \text{ iff }\ x_n=y_n\ \text{ for almost all }\ n.$$

Assume we are given a Γ-valued 1-cocycle c on $\mathcal{R}$. An important example is the Radon-Nikodym cocycle c_μ. It is the $\mathbb{R}^*_+$-valued cocycle defined by

$$c_\mu(x,y)=\prod_{n=1}^\infty\frac{\mu_n(y_n)}{\mu_n(x_n)}.$$

More generally, we say that a Γ-valued cocycle c is of product type if there exist maps $\theta_n\colon X_n\to\Gamma$ such that

$$c(x,y)=\prod_{n=1}^\infty\theta_n(y_n)\theta_n(x_n)^{-1}.$$

It is not difficult to see that a cocycle c is of product type if and only if it is constant on the sets

$$\{(x,y)\mid x_n=a,\ y_n=b \text{ and } x_m=y_m \text{ for } m\ne n\}\subset\mathcal{R}$$

for all $n\in\mathbb{N}$ and $a,b\in X_n$.

For a finite subset $I\subset\mathbb{N}$ and $a\in\prod_{n\in I}X_n$ put

$$Z(a):=\{x\in X\mid x_n=a_n \text{ for } n\in I\}.$$

Motivated by [1] define the asymptotic ratio set $r(c)$ of a Γ-valued cocycle c as the set of elements $\gamma\in\Gamma$ such that for every neighbourhood U of γ there exist numbers $t>s>0$, a sequence $\{I_n\}_{n=1}^\infty$ of mutually disjoint finite subsets of $\mathbb{N}$, subsets $K_n,L_n\subset\prod_{k\in I_n}X_k$ and bijections $T_n\colon K_n\to L_n$ such that

$$c_\mu(x,\bar T_nx)\in[s,t] \text{ and } c(x,\bar T_nx)\in U \text{ for all } x\in Z(a),\ a\in K_n \text{ and } n\ge1,$$

where $\bar T_n$ denotes the obvious map $\cup_{a\in K_n}Z(a)\to\cup_{b\in L_n}Z(b)$ defined by $T_n\colon K_n\to L_n$, and

$$\sum_{n=1}^\infty\sum_{a\in K_n}\mu(Z(a))=\infty.$$

Proposition 2.2. *For any Γ-valued 1-cocycle c on $\mathcal{R}$ we have $r(c)\subset r^*(c)$. If the cocycle c is of product type and for every $\gamma\in\Gamma$ there exist numbers $t>s>0$ and a neighbourhood U of γ such that $c_\mu(x,y)\in[s,t]$ as long as $c(x,y)\in U$, then $r(c)=r^*(c)$.*

Note that for the existence of s and t in the above proposition it suffices to have a continuous homomorphism $p\colon\Gamma\to\mathbb{R}^*_+$ such that $p\circ c=c_\mu$. Similarly, if such p exists, then the requirements on s, t and c_μ in the definition of the asymptotic ratio set can be omitted.

Proposition 2.2 is well-known for the Radon-Nikodym cocycle c_μ and is considered to be obvious in this case [9]. A rather detailed proof is contained in [17, Appendix A]. Essentially the same arguments work in the general case. Let us only sketch a proof of the inclusion $r(c)\subset r^*(c)$, which is the one we are going to use later.

Assume $\gamma\in r(c)$. Fix a neighbourhood U of γ. Choose s, t, I_n, K_n, L_n and T_n as in the definition of the asymptotic ratio set. Consider the sets $A_n:=\cup_{a\in K_n}Z(a)$ and $B_n:=\cup_{b\in L_n}Z(b)$, and put

$$A:=\cup_n\left(A_n\setminus\cup_{m<n}(A_m\cup B_m)\right)\ \text{ and }\ B:=\cup_n\left(B_n\setminus\cup_{m<n}(A_m\cup B_m)\right).$$

Since the sets $A_n\cup B_n$ are mutually independent and

$$\sum_n\mu(A_n\cup B_n)\ge\sum_n\mu(A_n)=\infty,$$

we have $\mu(A\cup B)=1$. Define a measurable bijective map $T\colon A\to B$ as follows: if

$$x\in(A_n\setminus\cup_{m<n}(A_m\cup B_m))\cap Z(a)$$

for some $a\in K_n$, then put $Tx:=\bar T_nx$. Then for any $x\in A$ we have $c_\mu(x,Tx)\in[s,t]$ and $c(x,Tx)\in U$.

Observe now that if we fixed a cylindrical set $Z(a)$ and then used only I_n, K_n, L_n and T_n with sufficiently large n in the above construction of A, B and T, then T would map $Z(a)\cap A$ onto $Z(a)\cap B$. We can therefore conclude that if $Z \subset X$ is a finite union of cylindrical sets $Z(a)$, then there exist measurable subsets $A, B \subset Z$ and a measurable bijective map $T\colon A \to B$ with graph in $\mathcal{R}$ such that

$$\mu(Z \setminus (A \cup B)) = 0,\ \ c_\mu(x, Tx) \in [s, t] \ \text{ and } \ c(x, Tx) \in U \ \text{ for } \ x \in A.$$

Any measurable subset $Z' \subset X$ of positive measure can be approximated arbitrarily well by such Z. Using that

$$\frac{dT^{-1}\mu}{d\mu}(x) = c_\mu(x, Tx) \in [s, t] \ \text{ for all } \ x \in A,$$

it is easy to see that if $\mu(Z'\Delta Z)$ is sufficiently small, then

$$\mu(Z' \cap T^{-1}(Z' \cap B)) > 0.$$

Therefore T maps the subset $Z' \cap T^{-1}(Z' \cap B)$ of Z' of positive measure into Z', and $c(x, Tx) \in U$ for any x in this subset. Since U was arbitrary, this shows that γ lies in the essential range of $c_{Z'}$.

3. DISTRIBUTION OF PRIME IDEALS IN NUMBER FIELDS

In this section we will prove Theorem 1.2 for number fields K. Set $n = [K : \mathbb{Q}] = r_1 + 2r_2$, so $\mathbb{A}_K$ and $\mathbb{A}_K^*$ each have $r_1 + r_2$ Archimedean places, with r_2 of them complex. Write $x := (x_v)_v$ for elements of $\mathbb{A}_K$, where v runs over the valuations of K. Valuations are normalized as in Tate's thesis [19, Sec. 2.1] and denoted $||\cdot||_v$ following Lang [12, XIV, §1], with normalized complex valuations being $||z||_{\mathbb{C}} := |z|^2$. The idelic norm map $N\colon \mathbb{A}_K^* \to \mathbb{R}_+^*$ sends $(x_v)_v \mapsto \prod_v ||x||_v$. For any non-zero integral ideal $\mathfrak{m}$, the quotient norm map $N_\mathfrak{m} : \mathbb{A}_K^*/K^*W_\mathfrak{m} \to \mathbb{R}_+^*$ is well-defined, and we put

$$\Gamma_\mathfrak{m} := \ker N_\mathfrak{m} \subset \mathbb{A}_K^*/K^*W_\mathfrak{m}.$$

The map $N\colon \mathbb{A}_K^* \to \mathbb{R}_+^*$ has a splitting homomorphism $s\colon \mathbb{R}_+^* \to \mathbb{A}_K^*$. Fix any such continuous homomorphism, for example, $s(t)_v = 1$ for $v \in V_f$ and $s(t) = (t^{1/n}, t^{1/n}, \dots, t^{1/n})$ at the $r_1 + r_2$ Archimedean places $v|\infty$ (note again that $||t^{1/n}||_v = t^{2/n}$ at complex places). Using this homomorphism we can identify $\mathbb{A}_K^*/K^*W_\mathfrak{m}$ with $\mathbb{R}_+^* \times \Gamma_\mathfrak{m}$. Then the embedding

$$I_\mathfrak{m} \hookrightarrow \mathbb{R}_+^* \times \Gamma_\mathfrak{m} = \mathbb{A}_K^*/K^*W_\mathfrak{m}$$

has the form $\mathfrak{a} \mapsto (N(\mathfrak{a})^{-1}, \rho_\mathfrak{m}(\mathfrak{a}))$ for some homomorphism $\rho_\mathfrak{m}\colon I_\mathfrak{m} \to \Gamma_\mathfrak{m}$. The group $\Gamma_\mathfrak{m}$ is a compact abelian Lie group whose connected components have real dimension $n - 1$. The points $\rho_\mathfrak{m}(\mathfrak{a}) \in \Gamma_\mathfrak{m}$ can be thought of as measuring *generalized angles* of ideals $\mathfrak{a} \in I_\mathfrak{m}$; see Sections 5.2 and 5.3 below.

Consider now the orbit equivalence relation defined by the action of $I_\mathfrak{m}$ on $\mathbb{A}_{S(\mathfrak{m})}/\hat{\mathcal{O}}^*_{S(\mathfrak{m})}$. This action is essentially free, so outside a set of measure zero we can define an $\mathbb{R}_+^* \times \Gamma_\mathfrak{m}$-valued 1-cocycle by

$$c(x, y) = (N(\mathfrak{a}), \rho_\mathfrak{m}(\mathfrak{a})) \ \text{ if } \ \mathfrak{a}x = y.$$

Furthermore, it is easy to see that the action of $I_\mathfrak{m}$ on $\mathbb{A}_{S(\mathfrak{m})}/\hat{\mathcal{O}}^*_{S(\mathfrak{m})}$ is ergodic. (Further justification is supplied in the proof of Theorem 1.2 below; more precisely, we will see there that the orbit equivalence relation on $\hat{\mathcal{O}}_{S(\mathfrak{m})}/\hat{\mathcal{O}}^*_{S(\mathfrak{m})}$ is ergodic, and since every $I_\mathfrak{m}$-orbit in $\mathbb{A}_{S(\mathfrak{m})}/\hat{\mathcal{O}}^*_{S(\mathfrak{m})}$ intersects $\hat{\mathcal{O}}_{S(\mathfrak{m})}/\hat{\mathcal{O}}^*_{S(\mathfrak{m})}$, it follows that the action of $I_\mathfrak{m}$ on $\mathbb{A}_{S(\mathfrak{m})}/\hat{\mathcal{O}}^*_{S(\mathfrak{m})}$ is also ergodic.) Therefore the statement of Theorem 1.2 is equivalent to the equality $r^*(c) = \mathbb{R}_+^* \times \Gamma_\mathfrak{m}$.

The computation of the asymptotic range of c will rely on the following distribution result.

Theorem 3.1. *The image under $\rho_\mathfrak{m}$ of the set of prime ideals in $\mathcal{O}$, ordered by the norm, is equidistributed in $\Gamma_\mathfrak{m}$. More precisely, for any continuous function f on $\Gamma_\mathfrak{m}$ we have*

$$\frac{1}{|\{\mathfrak{p} \notin S(\mathfrak{m}) : N(\mathfrak{p}) \le x\}|} \sum_{\mathfrak{p} \notin S(\mathfrak{m}) : N(\mathfrak{p}) \le x} f(\rho_\mathfrak{m}(\mathfrak{p})) \to \int_{\Gamma_\mathfrak{m}} f \, d\lambda_\mathfrak{m} \ \ \text{as} \ \ x \to \infty,$$

where $\lambda_\mathfrak{m}$ is the normalized Haar measure on the compact group $\Gamma_\mathfrak{m}$.

Proof. In order to prove the theorem it suffices to show that for any non-trivial character χ of $\Gamma_{\mathfrak{m}}$ we have

$$\frac{1}{|\{\mathfrak{p} \notin S(\mathfrak{m}) : N(\mathfrak{p}) \le x\}|} \sum_{\mathfrak{p} \notin S(\mathfrak{m}) : N(\mathfrak{p}) \le x} \chi(\rho_{\mathfrak{m}}(\mathfrak{p})) \to 0 \text{ as } x \to \infty.$$

In other words, since $|\{\mathfrak{p} : N(\mathfrak{p}) \le x\}| \sim x/\log x$ by the prime ideal theorem, we have to show

$$\sum_{\mathfrak{p} \notin S(\mathfrak{m}) : N(\mathfrak{p}) \le x} (\chi \circ \rho_{\mathfrak{m}})(\mathfrak{p}) = o\left(\frac{x}{\log x}\right). \tag{3.1}$$

But by construction of $\rho_{\mathfrak{m}}$, the character $\chi \circ \rho_{\mathfrak{m}}$ of $I_{\mathfrak{m}}$ is a Hecke character $(\bmod\ \mathfrak{m})$ that is not of the form $\mathfrak{a} \mapsto N(\mathfrak{a})^{it}$, $t \in \mathbb{R}$. For such characters (3.1) is well-known, see e.g. Narkiewicz [14, Prop. 7.17 (3rd Ed.); Prop. 7.9 (2nd Ed.)]. See also Section 5.2. below. ☐

Corollary 3.2. *For any subset $A \subset \Gamma_{\mathfrak{m}}$ with boundary of measure zero we have*

$$|\{\mathfrak{p} \notin S(\mathfrak{m}) : N(\mathfrak{p}) \le x \textit{ and } \rho_{\mathfrak{m}}(\mathfrak{p}) \in A\}| = \lambda_{\mathfrak{m}}(A)\frac{x}{\log x} + o\left(\frac{x}{\log x}\right) \textit{ as } x \to \infty.$$

Corollary 3.3. *For any subset $A \subset \Gamma_{\mathfrak{m}}$ with boundary of measure zero and any $\delta > 0$ we have*

$$|\{\mathfrak{p} \notin S(\mathfrak{m}) : x < N(\mathfrak{p}) \le (1+\delta)x \textit{ and } \rho_{\mathfrak{m}}(\mathfrak{p}) \in A\}| = \lambda_{\mathfrak{m}}(A)\delta\frac{x}{\log x} + o\left(\frac{x}{\log x}\right) \textit{ as } x \to \infty.$$

Proof of Theorem 1.2 for number fields. We are given an ideal $\mathfrak{m}$. Consider the subset $\hat{\mathcal{O}}_{S(\mathfrak{m})}/\hat{\mathcal{O}}^*_{S(\mathfrak{m})} \subset \mathbb{A}_{S(\mathfrak{m})}/\hat{\mathcal{O}}^*_{S(\mathfrak{m})}$, which is of measure one. Modulo a set of measure zero it can be identified with the product of spaces $\mathcal{O}_{\mathfrak{p}}^{\times}/\mathcal{O}_{\mathfrak{p}}^* \cong \mathbb{Z}_+$, for $\mathfrak{p} \notin S(\mathfrak{m})$, where $\mathbb{Z}_+ = \mathbb{Z}_{\ge 0}$ is the additive semigroup. Then the measure is the product of measures $\mu_{\mathfrak{p}}$ on $\mathbb{Z}_+$ defined by

$$\mu_{\mathfrak{p}}(n) := N(\mathfrak{p})^{-n}(1 - N(\mathfrak{p})^{-1}).$$

In this picture the orbit equivalence relation on $\hat{\mathcal{O}}_{S(\mathfrak{m})}/\hat{\mathcal{O}}^*_{S(\mathfrak{m})}$ is exactly the ergodic equivalence relation considered in the previous section:

$$a \sim b \text{ if and only if } a_{\mathfrak{p}} = b_{\mathfrak{p}} \text{ for almost all } \mathfrak{p},$$

and the cocycle $c_{\hat{\mathcal{O}}_{S(\mathfrak{m})}/\hat{\mathcal{O}}^*_{S(\mathfrak{m})}}$ is the cocycle of product type defined by the maps $\theta_{\mathfrak{p}} \colon \mathbb{Z}_+ \to \mathbb{R}^*_+ \times \Gamma_{\mathfrak{m}}$,

$$\theta_{\mathfrak{p}}(n) := (N(\mathfrak{p})^n, \rho_{\mathfrak{m}}(\mathfrak{p})^n).$$

Note that these formulas show in particular that if we define a homomorphism $p \colon \mathbb{R}^*_+ \times \Gamma_{\mathfrak{m}} \to \mathbb{R}^*_+$ by $p(x,\gamma) := x^{-1}$, then $p \circ c$ coincides with the Radon-Nikodym cocycle.

In order to prove the theorem, by Proposition 2.1 it suffices to show that the asymptotic range of $c_{\hat{\mathcal{O}}_{S(\mathfrak{m})}/\hat{\mathcal{O}}^*_{S(\mathfrak{m})}}$ is equal to $\mathbb{R}^*_+ \times \Gamma_{\mathfrak{m}}$. Fix a point $(x_0, y_0) \in \mathbb{R}^*_+ \times \Gamma_{\mathfrak{m}}$. Since the asymptotic range is a closed subgroup, it suffices to consider $x_0 > 1$. By Proposition 2.2 it is enough to show that (x_0, y_0) belongs to the asymptotic ratio set of $c_{\hat{\mathcal{O}}_{S(\mathfrak{m})}/\hat{\mathcal{O}}^*_{S(\mathfrak{m})}}$. This is done along familiar lines [2, 16], as follows.

Fix a neighbourhood U of (x_0, y_0). Choose $\varepsilon > 0$ and an open subset $V \subset \Gamma_{\mathfrak{m}}$ with boundary of measure zero such that

$$(x_0 - \varepsilon, x_0 + \varepsilon) \times y_0 V V^{-1} \subset U.$$

Choose $\delta > 0$ such that $1 + \delta < x_0$ and $\delta x_0 < \varepsilon$. Define sets B_n of prime ideals by

$$B_{2k} = \{\mathfrak{p} \notin S(\mathfrak{m}) : x_0^{2k} < N(\mathfrak{p}) \le (1+\delta)x_0^{2k} \text{ and } \rho_{\mathfrak{m}}(\mathfrak{p}) \in V\},$$

$$B_{2k+1} = \{\mathfrak{p} \notin S(\mathfrak{m}) : x_0^{2k+1} < N(\mathfrak{p}) \le (1+\delta)x_0^{2k+1} \text{ and } \rho_{\mathfrak{m}}(\mathfrak{p}) \in y_0 V\}.$$

By the choice of δ the sets B_n are pairwise disjoint. By Corollary 3.3 we have

$$|B_n| \sim \frac{\lambda_{\mathfrak{m}}(V)\delta x_0^n}{n \log x_0} \text{ as } n \to \infty.$$

In particular, there exists k_0 such that $|B_{2k+1}| \ge |B_{2k}|$ for $k \ge k_0$. For every $k \ge k_0$ choose a subset $C_{2k+1} \subset B_{2k+1}$ such that $|C_{2k+1}| = |B_{2k}|$. Let $\mathfrak{p}_1, \mathfrak{p}_2, \dots$ be the elements of $\cup_{k\ge k_0} B_{2k}$ enumerated so that $N(\mathfrak{p}_1) \le N(\mathfrak{p}_2) \le \dots$, and let $\mathfrak{q}_1, \mathfrak{q}_2, \dots$ be the elements of $\cup_{k\ge k_0} C_{2k+1}$ enumerated in the same way. Then, if $\mathfrak{p}_n \in B_{2k}$ for some k, we have $\mathfrak{q}_n \in B_{2k+1}$, and therefore

$$N(\mathfrak{q}_n\mathfrak{p}_n^{-1}) \in (x_0 - \varepsilon, x_0 + \varepsilon) \text{ and } \rho_{\mathfrak{m}}(\mathfrak{q}_n\mathfrak{p}_n^{-1}) \in y_0VV^{-1}.$$

We also have

$$\sum_{n=1}^{\infty} N(\mathfrak{p}_n)^{-1} \ge \sum_{k=k_0}^{\infty} \frac{|B_{2k}|}{(1+\delta)x_0^{2k}} = \infty.$$

Hence for the sets I_n, K_n and L_n required by the definition of the asymptotic ratio set we can take $I_n = \{\mathfrak{p}_n, \mathfrak{q}_n\}$, $K_n = \{(1,0)\} \subset \mathbb{Z}_+^{I_n}$ and $L_n = \{(0,1)\}$. □

4. FUNCTION FIELDS

In this section we will prove Theorem 1.2 for function fields. The strategy is the same as for number fields, but some changes are necessary, since for function fields the Chebotarev density theorem does not hold for the natural density (the density oscillates as a function of x with no limiting value), and therefore the straightforward analogue of Theorem 3.1 cannot be true.

Assume K is a global function field with constant field $\mathbb{F}_q$. In this case the image of the norm map $N\colon \mathbb{A}_K^* \to \mathbb{R}_+^*$ is $q^{\mathbb{Z}}$. Fix an idele a of norm q. Then for a given nonzero ideal $\mathfrak{m} \subset \mathcal{O}$ the group $\mathbb{A}_K^*/K^*W_{\mathfrak{m}}$ can be identified with $a^{\mathbb{Z}} \times \Gamma_{\mathfrak{m}} \cong \mathbb{Z} \times \Gamma_{\mathfrak{m}}$, where as before $\Gamma_{\mathfrak{m}}$ is the kernel of $N_{\mathfrak{m}}\colon \mathbb{A}_K^*/K^*W_{\mathfrak{m}} \to q^{\mathbb{Z}}$. We have to show that the diagonal action of $I_{\mathfrak{m}}$ on $\mathbb{Z} \times \Gamma_{\mathfrak{m}} \times \mathbb{A}_{S(\mathfrak{m})}/\hat{\mathcal{O}}_{S(\mathfrak{m})}^*$ is ergodic. Since the group $\mathcal{O}_\infty^*$ is profinite, we can make one more reduction and replace $\Gamma_{\mathfrak{m}}$ by the quotient by a compact open subgroup $U \subset \mathcal{O}_\infty^*$. The group $\Gamma_{\mathfrak{m}}/U$ is finite, and it is canonically isomorphic to the quotient of $\mathbb{A}_K^*$ by the open subgroup $a^{\mathbb{Z}}K^*UW_{\mathfrak{m}}$. By class field theory it follows that $\Gamma_{\mathfrak{m}}/U$ is the Galois group of a finite abelian extension L/K.

Lemma 4.1. *The extension L/K is geometric; that is, K and L have the same field of constants. It is unramified at all finite places $\mathfrak{p} \notin S(\mathfrak{m})$.*

Proof. Assume the constant field of L is $\mathbb{F}_{q^m}$. The Artin map $\mathbb{A}_K^* \to \mathrm{Gal}(L/K)$ defines a homomorphism $\mathbb{A}_K^* \to \mathrm{Gal}(\mathbb{F}_{q^m}/\mathbb{F}_q)$. This homomorphism factors through the divisor group of K and maps a divisor D to the Frobenius raised to the power $\deg D$. But by definition of L our fixed idele a acts trivially on L, so the corresponding divisor of degree 1 acts trivially on $\mathbb{F}_{q^m}$. Hence $m = 1$, so the extension is geometric. Since $\hat{\mathcal{O}}_{S(\mathfrak{m})}^*$ is contained in the kernel of the homomorphism $\mathbb{A}_K^* \to \mathrm{Gal}(L/K)$, the extension is unramified at all finite places $\mathfrak{p} \notin S(\mathfrak{m})$. □

Therefore in order to prove Theorem 1.2 for function fields it suffices to establish the following.

Theorem 4.2. *For any non-zero ideal $\mathfrak{m} \subset \mathcal{O}$ and any finite abelian geometric extension L/K unramified at all finite places $\mathfrak{p} \notin S(\mathfrak{m})$, the action of $I_{\mathfrak{m}}$ on $\mathbb{Z} \times \mathrm{Gal}(L/K) \times \mathbb{A}_{S(\mathfrak{m})}/\hat{\mathcal{O}}_{S(\mathfrak{m})}^*$, given by*

$$\mathfrak{a}(n, g, x) = (n + \deg \mathfrak{a}, \sigma(\mathfrak{a})g, \mathfrak{a}x),$$

where $\sigma\colon I_{\mathfrak{m}} \to \mathrm{Gal}(L/K)$ is the Artin map, is ergodic.

This result is, in fact, a consequence of [17, Theorem 2.1], but here we will sketch a slightly different proof following the arguments of the previous section. It should also be remarked that the theorem is not true for non-geometric extensions, see Section 5.5.

Proof of Theorem 4.2. Similarly to the proof of Theorem 1.2 for number fields define a $\mathbb{Z}\times\mathrm{Gal}(L/K)$-valued cocycle c on the orbit equivalence relation on $\mathbb{A}_{S(\mathfrak{m})}/\hat{\mathcal{O}}^*_{S(\mathfrak{m})}$ by

$$c(x,y)=(\deg\mathfrak{a},\sigma(\mathfrak{a}))\ \text{ if }\ \mathfrak{a}x=y.$$

We aim to show that the asymptotic range of c is $\mathbb{Z}\times\mathrm{Gal}(L/K)$. For this it suffices to show that for any $g\in\mathrm{Gal}(L/K)$ the element $(1,g)$ belongs to the asymptotic ratio set of the restriction of c to the equivalence relation on $\prod_{\mathfrak{p}\notin S(\mathfrak{m})}\mathcal{O}_{\mathfrak{p}}^\times/\mathcal{O}_{\mathfrak{p}}^*\cong\prod_{\mathfrak{p}\notin S(\mathfrak{m})}\mathbb{Z}_+$. For this, define sets B_n of prime ideals by

$$B_{2k}=\{\mathfrak{p}\notin S(\mathfrak{m}):\deg\mathfrak{p}=2k\text{ and }\sigma(\mathfrak{p})=e\},$$

$$B_{2k+1}=\{\mathfrak{p}\notin S(\mathfrak{m}):\deg\mathfrak{p}=2k+1\text{ and }\sigma(\mathfrak{p})=g\}.$$

By the Chebotarev density theorem for function fields, see e.g. [6, Proposition 6.4.8], we have

$$|B_n|=\frac{q^n}{n[L:K]}+O(q^{n/2})\ \text{ as }\ n\to\infty.$$

Proceeding now as in the proof of Theorem 1.2 for number fields, we conclude that $(1,g)$ belongs to the asymptotic ratio set. □

5. REMARKS

5.1. Ergodicity of an action implies that the orbit of almost every point is dense. However the converse assertion is not true in general. Here it is not difficult to show directly the density of almost every orbit of the action of K^* on $\mathbb{A}_K$. More precisely, the orbit of every non-invertible adele a such that $a_v\ne0$ for all places v is dense. This was observed for $K=\mathbb{Q}$ by Laca and Raeburn [11, Lemma 3.2]. In order to prove the same for an arbitrary global field K fix $b\in\mathbb{A}_K$ and a neighbourhood U of b. We want to find $x\in K^*$ such that $xa\in U$. The set U contains an open subset of the form $V\times(b_f+\hat{\mathfrak{a}})$, where $V\subset K_\infty\setminus\{0\}$, $\mathfrak{a}$ is a non-zero ideal in $\mathcal{O}$ and $\hat{\mathfrak{a}}$ is the closure of $\mathfrak{a}$ in $\hat{\mathcal{O}}$.

We claim that there exists $q\in K^*$ such that $qa_f\in\hat{\mathfrak{a}}$ and $a_\infty^{-1}V+q\mathcal{O}=K_\infty$. In order to see this choose $y\in\mathcal{O}^\times$ such that $ya_f\in\hat{\mathfrak{a}}$. Consider the set $\{v_1,v_2,\dots\}$ of finite places $v\notin S(\mathfrak{a})$ such that $\|a_v\|_v<1$. Note that since a is not invertible and $a_v\ne0$ for all v, this set is infinite. Consider the sequence of ideals $\mathfrak{m}_n=\mathfrak{p}_{v_1}\dots\mathfrak{p}_{v_n}$ in $\mathcal{O}$. Choose a subsequence $\{\mathfrak{m}_{n_k}\}_{k=1}^\infty$ belonging to the same ideal class. Then the ideals $\mathfrak{m}_{n_1}^{-1}\mathfrak{m}_{n_k}=\mathfrak{p}_{v_{n_1+1}}\mathfrak{p}_{v_{n_1+2}}\dots\mathfrak{p}_{v_{n_k}}$ are principal. Let z_k be a generator of $\mathfrak{m}_{n_1}^{-1}\mathfrak{m}_{n_k}$. Then the elements $q_k=z_k^{-1}y\in K^*$ have the properties that $q_ka_f\in\hat{\mathfrak{a}}$ and $N((q_k))\to0$ as $k\to\infty$. Since $\mathcal{O}$ is a cocompact lattice in K_∞ and since for appropriate units $u_k\in\mathcal{O}^*$ we have $\|u_kq_k\|_v\to0$ for every $v|\infty$, for sufficiently large k we get $a_\infty^{-1}V+q_k\mathcal{O}=K_\infty$. Thus our claim is proved.

We can now find the required element x. First, using density of K in $\mathbb{A}_{K,f}$, choose $x'\in K$ such that $x'a_f\in b_f+\hat{\mathfrak{a}}$. Next, using that $a_\infty^{-1}V+q\mathcal{O}=K_\infty$, choose $y\in\mathcal{O}$ such that $x'+qy\in a_\infty^{-1}V$. Then the element $x=x'+qy\in K$ has the properties that $xa_\infty\in V$ and $xa_f=x'a_f+qya_f\in b_f+\hat{\mathfrak{a}}$, as $qa_f\in\hat{\mathfrak{a}}$. Therefore $xa\in V\times(b_f+\hat{\mathfrak{a}})\subset U$. Since $0\notin V$, we must have $x\in K^*$.

5.2. For general number fields K the key result underlying the proof of Theorem 1.1 is Theorem 3.1, which gives equidistribution over the group $\Gamma_{\mathfrak{m}}$. This result uses the full set of Hecke characters (“Grössencharacters”) of the field K, as introduced in Hecke [7], [8], and given in Narkiewicz [14, Sect. 7.3 (2nd Ed)] or Neukirch [18, Chap. VII.6]. If $K\ne\mathbb{Q}$ then the set of such Grössencharacters includes characters of infinite order. The equidistribution result in Theorem 3.1 for number fields is at bottom derived using the fact that the L-functions of Grössencharacters are free from zeros and poles on the line $\mathrm{Re}(s)=1$, except for the characters $N(\cdot)^{it}$ having simple poles at $s=1-it$. Since $L(s,N(\cdot)^{it}\chi)=L(s+it,\chi)$, it makes sense to normalize characters, that is, to fix representatives among the characters that differ by a factor $N(\cdot)^{it}$. In this and the next subsection we fix the section $s(t)=(t^{1/n},\dots,t^{1/n},1,1,\dots)$ of the norm map, and call a character of $\mathbb{A}_K^*/K^*$ normalized if it is trivial on $s(\mathbb{R}_+^*)$. This normalization agrees with Lang [12], but differs from Narkiewicz [14].

In order to describe the nature of the groups $\mathbb{A}_K/K^*W_\mathfrak{m}$ and $\Gamma_\mathfrak{m}$ in more classical language, observe that the quotient of $\mathbb{A}_K/K^*W_\mathfrak{m}$ by K^*_∞ is isomorphic to the finite group $C_\mathfrak{m} = I_\mathfrak{m}/\mathcal{P}_{\mathfrak{m},1}$, the ray class group modulo $\mathfrak{m}$. We therefore have a short exact sequence

$$1 \to K^*_\infty/U_{\mathfrak{m},1} \to \mathbb{A}_K/K^*W_\mathfrak{m} \to C_\mathfrak{m} \to 1.$$

Hence $\mathbb{A}_K/K^*W_\mathfrak{m}$ is an abelian Lie group with finitely many connected components, and with the connected component of the identity isomorphic to $K^+_\infty/U^+_{\mathfrak{m},1}$, where K^+_∞ is the connected component of the identity in K^*_∞ and $U^+_{\mathfrak{m},1} = U_{\mathfrak{m},1} \cap K^+_\infty$ is the group of totally positive units $\bmod\ \mathfrak{m}$.

For an Archimedean place v put $N_v = 1$ if v is real and $N_v = 2$ if v is complex. Denote by $K^*_{\infty,1} \subset K^*_\infty$ the subgroup of elements x such that $\prod_{v|\infty} ||x_v||_v = \prod_{v|\infty} |x_v|^{N_v} = 1$. Then we also have a short exact sequence

$$1 \to K^*_{\infty,1}/U_{\mathfrak{m},1} \to \Gamma_\mathfrak{m} \to C_\mathfrak{m} \to 1.$$

The homomorphism $\rho_\mathfrak{m}\colon I_\mathfrak{m} \to \Gamma_\mathfrak{m}$ is such that the image of $\rho_\mathfrak{m}(\mathfrak{a})$ in $C_\mathfrak{m}$ is the class $[\mathfrak{a}]$ of $\mathfrak{a}$, and if $\mathfrak{a} = (x)$ for some $x \in K^*_{\mathfrak{m},1}$, then

$$\rho_\mathfrak{m}(\mathfrak{a}) = s_\infty(N(\mathfrak{a}))x^{-1}U_{\mathfrak{m},1} \in K^*_{\infty,1}/U_{\mathfrak{m},1} \subset \Gamma_\mathfrak{m},$$

where $s_\infty(t) = (t^{1/n}, \dots, t^{1/n}) \in K^*_\infty$. More explicitly, the group $\Gamma_\mathfrak{m}$ can be described as the quotient of $I_\mathfrak{m} \times K^*_{\infty,1}/U_{\mathfrak{m},1}$ by $\mathcal{P}_{\mathfrak{m},1}$ under the diagonal embedding, where the homomorphism $\mathcal{P}_{\mathfrak{m},1} \to K^*_{\infty,1}/U_{\mathfrak{m},1}$ is given by $(x) \mapsto s_\infty(N((x)))^{-1}xU_{\mathfrak{m},1}$ for $x \in K^*_{\mathfrak{m},1}$.

Let r_1 and r_2 be the number of real and complex places of K, respectively, with $n = r_1 + 2r_2$. Then $K_\infty = \mathbb{R}^{r_1} \times \mathbb{C}^{r_2}$. The connected component $K^+_\infty/U^+_{\mathfrak{m},1}$ of the identity in $\mathbb{A}_K/K^*W_\mathfrak{m}$ has the form $\mathbb{R}^{r_1+2r_2}/\Lambda_\mathfrak{m}$, for a certain lattice $\Lambda_\mathfrak{m}$ of (non-full) rank $r_1 + 2r_2 - 1$ which (under a logarithmic change of variable) encodes the unit lattice together with additional lattice vectors encoding the ambiguity of the argument at each complex place. The connected component of the identity of the smaller group $\Gamma_\mathfrak{m}$ is then $V/\Lambda_\mathfrak{m}$, where V is the real subspace of $\mathbb{R}^n$ of dimension $n-1$ spanned by the lattice $\Lambda_\mathfrak{m}$. We emphasize that the lattice Λ_m has rank $n-1$ which for $r_2 > 0$ is larger than the standard logarithmic encoding in $\mathbb{R}^{r_1+r_2}$ of the unit lattice of rank $r_1 + r_2 - 1$ used for computing the regulator of the field K (as in Lang [12, Chap. V]). The standard encoding detects only size information on the units, while the lattice $\Lambda_\mathfrak{m}$ contains an extra r_2 components that encode information on the complex arguments ("angles") of the units.

We give an example. Take $K = \mathbb{Q}(\theta)$ to be a (non-Galois) real cubic field of discriminant $d_K = -23$ obtained by adjoining the real root of $X^3 - X - 1 = 0$. This equation has one real root $\theta \approx 1.3247$ and two conjugate complex roots $\vartheta^\pm = \frac{1}{\sqrt{\theta}} e^{\pm 2\pi i \phi}$. An important feature is that the angle ϕ is irrational, because all nonzero powers of ϑ^+ are non-real, since $\mathbb{Q}(\vartheta^+)$ contains no real units except ± 1. It is known that the field K has ring of integers $\mathcal{O} = \mathbb{Z}[1, \theta, \theta^2]$. The unit rank of K is $r_1 + r_2 - 1 = 1$, and θ is a fundamental unit of K. We choose $\mathfrak{m} = (1)$ to be the unit ideal. The field K is known to have class number 1, so that $I_\mathfrak{m} = \mathcal{P}_{\mathfrak{m},1}$, and the group $\mathbb{A}^*_K/K^*W_\mathfrak{m} = \mathbb{A}^*_K/K^*\hat{\mathcal{O}}^*$ is isomorphic to $K^*_\infty/U_{\mathfrak{m},1}$, with $U_{\mathfrak{m},1} = U_K$ being the full unit group. Since $-1 \in U_{\mathfrak{m},1}$, this group is connected, so it coincides with K^+_∞/U^+_K, where U^+_K is the group of totally positive units, and $K^+_\infty = \mathbb{R}^*_+ \times \mathbb{C}^*$. Under the logarithmic change of coordinate sending $\log\colon \mathbb{R}^*_+ \to \mathbb{R}$ and $\log\colon \mathbb{C}^* \to \mathbb{R} \oplus \mathbb{R}/2\pi\mathbb{Z}$, the group U^+_K is mapped to a 1-dimensional lattice in $\mathbb{R} \oplus \mathbb{R} \oplus \mathbb{R}/2\pi\mathbb{Z}$, with basis vector:

$$\mathbf{v}_1 = (\log\theta, -\frac{1}{2}\log\theta, 2\pi\phi),$$

with the argument defined $(\bmod\ 2\pi)$. We can lift this lattice to $\mathbb{R}^3$, to obtain a 2-dimensional lattice with a second basis vector coming from the ambiguity of the argument:

$$\mathbf{v}_2 = (0, 0, 2\pi).$$

Thus we obtain a 2-dimensional lattice $\Lambda_K := \mathbb{Z}[\mathbf{v}_1, \mathbf{v}_2]$ inside $\mathbb{R}^3$. Then the group $\mathbb{A}^*_K/K^*\hat{\mathcal{O}}^* = K^+_\infty/U^+_K$ can be identified under the logarithmic map with $\mathbb{R}^3/\Lambda_K$, with $\mathbf{v}_2$ removing the ambiguity of the branch of the logarithm. Now let V be the 2-dimensional subspace $\mathbb{R}[\mathbf{v}_1, \mathbf{v}_2]$, given explicitly by $V = \{\mathbf{x} = (x_1, x_2, x_3) : x_1 + 2x_2 = 0\}$. Then we may similarly identify $\Gamma_\mathfrak{m} = \Gamma_1$ with V/Λ_K, which is a torus of dimension

$n-1=2$. The normalized Grössencharacters $(\mathrm{mod}\ 1)$ form the dual of Γ_1, that is, a group isomorphic to $\mathbb{Z}^2$; all but one of these are characters of infinite order. These characters take the angles in the complex embeddings of K into account.

In order to give these Grössencharacters explicitly we need a formula for the projection $\mathbb{R}^3 \to V$ along the line $\mathbb{R}(1,1,0) \subset \mathbb{R}^3$, which corresponds to our fixed embedding of $\mathbb{R}^*_+$ into $\mathbb{A}^*_K$. For this, take the dual basis $\{\mathbf{w}_1, \mathbf{w}_2\}$ to $\{\mathbf{v}_1, \mathbf{v}_2\}$ viewed in the subspace W orthogonal to $\mathbb{R}(1,1,0)$ with respect to the standard scalar product $\langle \mathbf{w}, \mathbf{x}\rangle = w_1x_1 + w_2x_2 + w_3x_3$:

$$\mathbf{w}_1 = (\frac{2}{3\log\theta}, -\frac{2}{3\log\theta}, 0),\ \ \mathbf{w}_2 = (-\frac{2\phi}{3\log\theta}, \frac{2\phi}{3\log\theta}, \frac{1}{2\pi}).$$

Then the projection is given by $\mathbf{x} \mapsto \langle \mathbf{w}_1, \mathbf{x}\rangle \mathbf{v}_1 + \langle \mathbf{w}_2, \mathbf{x}\rangle \mathbf{v}_2$. Now, given a non-zero ideal $\mathfrak{a} = (\alpha)$ in K, by multiplying by -1 if necessary we may suppose that the generator $\alpha > 0$. Then the algebraic conjugate of α in the field $\mathbb{Q}(\vartheta^+)$ takes the form $\alpha^+ = \sqrt{\frac{N(\mathfrak{a})}{\alpha}} e^{2\pi i\psi}$, where $\psi = \psi(\alpha)$ is well-defined $(\mathrm{mod}\, 1)$. The normalzied Grössencharacter χ_{k_1,k_2} corresponding to $(k_1,k_2) \in \mathbb{Z}^2$ assigns $\mathfrak{a}$ the value

$$\chi_{k_1,k_2}(\mathfrak{a}) := e^{-2\pi i(k_1\langle \mathbf{w}_1,\mathbf{x}\rangle + k_2\langle \mathbf{w}_2,\mathbf{x}\rangle)},$$

where $\mathbf{x} = \mathbf{x}(\alpha) = (\log\alpha, -\frac{1}{2}\log\alpha + \frac{1}{2}\log N(\mathfrak{a}), 2\pi\psi)$ is the image of α under the logarithm map.

5.3. Theorem 3.1, or rather Corollary 3.2, generalizes a result of Mitsui [13, Section 3] on distribution of prime ideal numbers, which in turn generalizes earlier results of Hecke and Rademacher for quadratic number fields. In order to see this consider the homomorphism $\theta\colon K^*_\infty \to K^+_\infty$ that replaces every coordinate x_v of $x \in K^*_\infty$ for a real Archimedean place v by $|x_v|$. Since the group $K^+_\infty/\theta(U_{\mathfrak{m},1})$ is divisible, the homomorphism $\theta\colon K^*_\infty/U_{\mathfrak{m},1} \to K^+_\infty/\theta(U_{\mathfrak{m},1})$ extends to a homomorphism $\theta_{\mathfrak{m}}\colon \mathbb{A}^*_K/K^*W_{\mathfrak{m}} \to K^+_\infty/\theta(U_{\mathfrak{m},1})$. The homomorphism $\theta_{\mathfrak{m}}$ maps $\Gamma_{\mathfrak{m}}$ onto $K^+_{\infty,1}/\theta(U_{\mathfrak{m},1})$, where $K^+_{\infty,1} = K^+_\infty \cap K^*_{\infty,1}$. It follows that the image of the prime ideals under $\theta_{\mathfrak{m}} \circ \rho_{\mathfrak{m}}$, ordered by the norm, is equidistributed in $K^+_{\infty,1}/\theta(U_{\mathfrak{m},1})$. But we can conclude more.

Fix an ideal $\mathfrak{a} \in I_{\mathfrak{m}}$ and consider its class $[\mathfrak{a}]$ in the ray class group $C_{\mathfrak{m}}$. Identifying the set of cosets of $K^*_{\infty,1}/U_{\mathfrak{m},1}$ in $\Gamma_{\mathfrak{m}}$ with $C_{\mathfrak{m}}$, by Theorem 3.1 we conclude that the prime ideals in the class $[\mathfrak{a}]$ have density $|C_{\mathfrak{m}}|^{-1} = h_K^{-1}\varphi(\mathfrak{m})^{-1}$ in the set of all prime ideals, and their image under $\rho_{\mathfrak{m}}$, ordered by the norm, is equidistributed in the corresponding coset of $K^*_{\infty,1}/U_{\mathfrak{m},1}$. The homomorphism $\theta_{\mathfrak{m}}$ maps surjectively every such coset onto $K^+_{\infty,1}/\theta(U_{\mathfrak{m},1})$. It follows that: *the set* $\{(\theta_{\mathfrak{m}} \circ \rho_{\mathfrak{m}})(\mathfrak{p}) \mid \mathfrak{p} \in [\mathfrak{a}]\}$*, ordered by the norm, is equidistributed in* $K^+_{\infty,1}/\theta(U_{\mathfrak{m},1})$.

Consider now *ideal numbers* $\hat{K}^* \subset K^*_{\mathbb{C}}$, where $K_{\mathbb{C}} = \prod_{v|\infty}\mathbb{C}$. They are constructed as follows (see e.g. [18, p. 486]). Let $\mathcal{P}$ be the group of principal fractional ideals, and I be the group of all fractional ideals. Since the group $K^*_{\mathbb{C}}$ is divisible, the canonical homomorphism $\mathcal{P} = K^*/\mathcal{O}^* \to K^*_{\mathbb{C}}/\mathcal{O}^*$ extends to a homomorphism $I \to K^*_{\mathbb{C}}/\mathcal{O}^*$. Consider the image of this homomorphism. Then take $\hat{K}^*$ to be its preimage in $K^*_{\mathbb{C}}$. The point of introducing ideal numbers is to provide a mechanism to assign "angles" to non-principal ideals, since elements of $K^*_{\mathbb{C}}$ have well-defined angles componentwise.

Fix an ideal number $\hat{a} \in \hat{K}^*$ representing $\mathfrak{a}$. Then every ideal $\mathfrak{b} \in I_{\mathfrak{m}}$ in the class $[\mathfrak{a}] \in C_{\mathfrak{m}}$ can be represented by a unique, up to a factor from $U_{\mathfrak{m},1}$, ideal number $\hat{b}$ lying in $K^*_{\mathfrak{m},1}\hat{a}$; in other words, $\hat{b} \in \hat{K}^*$ is such that $\hat{b} \equiv \hat{a} \bmod \mathfrak{m}$. The homomorphism $\theta\colon K^*_\infty \to K^+_\infty$ constructed earlier has an extension to $K^*_{\mathbb{C}}$ defined in the same way. We therefore get an element $\theta(\hat{b})\theta(U_{\mathfrak{m},1}) \in K^+_\infty/\theta(U_{\mathfrak{m},1})$ that depends on $\mathfrak{b}$ and $\hat{a}$, but not on the choice of $\hat{b}$. Finally, define $\rho(\mathfrak{b}) := s_\infty(N(\mathfrak{b}))^{-1}\theta(\hat{b})\theta(U_{\mathfrak{m},1}) \in K^+_{\infty,1}/\theta(U_{\mathfrak{m},1})$. Then the result of Mitsui can be formulated by saying that for certain sets $A \subset K^+_{\infty,1}/\theta(U_{\mathfrak{m},1})$ with boundary of measure zero we have

$$|\{\mathfrak{p} \in [\mathfrak{a}] \mid N(\mathfrak{p}) \le x \text{ and } \rho(\mathfrak{p}) \in A\}| = \frac{\tilde{\lambda}_{\mathfrak{m}}(A)x}{h_K\varphi(\mathfrak{m})\log x} + o\left(\frac{x}{\log x}\right),$$

where $\tilde{\lambda}_{\mathfrak{m}}$ is the normalized Haar measure on $K^+_{\infty,1}/\theta(U_{\mathfrak{m},1})$. But by construction we have

$$\rho(\mathfrak{p}) = \rho(\mathfrak{a})(\theta_{\mathfrak{m}} \circ \rho_{\mathfrak{m}})(\mathfrak{a})(\theta_{\mathfrak{m}} \circ \rho_{\mathfrak{m}})(\mathfrak{p})^{-1},$$

so this is the same equidistribution result as the one stated in italics above. Mitsui's result, however, is stronger in that he obtains an estimate of the remainder term for sets A of a particular shape, and obtaining a remainder term is where the main work of Mitsui's paper lies.

5.4. As was already mentioned in the introduction, our ergodicity result is closely related to the analysis of Bost-Connes type systems. To make the connection more precise let us state the result in yet another form. Let $C_K = \mathbb{A}^*_K/K^*$ be the idele class group, and consider the balanced product $C_K \times_{\hat{\mathcal{O}}^*} \mathbb{A}_{K,f}$. The action $g(x, y) = (xg^{-1}, gy)$ of $\mathbb{A}^*_{K,f}$ on $\mathbb{A}^*_K \times \mathbb{A}_{K,f}$ defines an action of the group $I \cong \mathbb{A}^*_{K,f}/\hat{\mathcal{O}}^*$ of fractional ideals on $C_K \times_{\hat{\mathcal{O}}^*} \mathbb{A}_{K,f}$. Then a statement equivalent to Theorem 1.1 is that the action of I on $C_K \times_{\hat{\mathcal{O}}^*} \mathbb{A}_{K,f}$ is ergodic. Indeed, as in Section 1 ergodicity of this action is equivalent to ergodicity of the action of I on $C_K \times_{\hat{\mathcal{O}}^*} (\mathbb{A}_{K,f}/W_{\mathfrak{m}}) = (C_K/W_{\mathfrak{m}}) \times_{\hat{\mathcal{O}}^*} \mathbb{A}_{K,f}$ for all $\mathfrak{m}$, which, in turn, is the same as ergodicity of the action of $I_{\mathfrak{m}}$ on $(C_K/W_{\mathfrak{m}}) \times (\mathbb{A}_{S(\mathfrak{m})}/\hat{\mathcal{O}}^*_{S(\mathfrak{m})})$, and this is exactly Theorem 1.2.

Assume now that K is a number field and consider the compact group $C^1_K = \ker(N \colon C_K \to \mathbb{R}^*_+)$. Identifying it with the quotient $C_K/s(\mathbb{R}^*_+)$, where s is as before a section of the norm map, we get an action of I on $C^1_K \times_{\hat{\mathcal{O}}^*} \mathbb{A}_{K,f}$. Then ergodicity of the action of I on $C_K \times_{\hat{\mathcal{O}}^*} \mathbb{A}_{K,f}$ means the following: *the action of I on $C^1_K \times_{\hat{\mathcal{O}}^*} \mathbb{A}_{K,f}$ is ergodic of type* III_1. For $K = \mathbb{Q}$ this is known to be equivalent to the statement of the Bost-Connes phase transition theorem [3] for the inverse temperature $\beta = 1$, see Remarks in [15]. For general number fields this is stronger than what has been proved for Bost-Connes type systems [10, 16], as in such systems the group C^1_K is replaced by its quotient $\mathrm{Gal}(K^{ab}/K) \cong C_K/\overline{K^+_\infty}$. The same arguments can be used for all $\beta \in (0, 1]$, and together with the technique developed in [10] for $\beta > 1$ this leads to the following generalization of the Bost-Connes phase transition theorem, formulated in dynamical systems terms.

Theorem. *For a number field K the following hold:*

(i) *for every $\beta \in (0, 1]$ there exists a unique Borel measure μ_β on $C^1_K \times_{\hat{\mathcal{O}}^*} \mathbb{A}_{K,f}$ with $\mu_\beta(C^1_K \times_{\hat{\mathcal{O}}^*} \hat{\mathcal{O}}) = 1$ and with $\mu_\beta(\mathfrak{a}\,\cdot) = N(\mathfrak{a})^{-\beta}\mu_\beta$ for all $\mathfrak{a} \in I$; furthermore, the action of I on $(C^1_K \times_{\hat{\mathcal{O}}^*} \mathbb{A}_{K,f}, \mu_\beta)$ is ergodic of type* III_1*;*

(ii) *for every $\beta \in (1, +\infty)$ there exists a bijective correspondence between points of $C^1_K \times_{\hat{\mathcal{O}}^*} \hat{\mathcal{O}}^* \cong C^1_K$ and extremal Borel measures ν on $C^1_K \times_{\hat{\mathcal{O}}^*} \mathbb{A}_{K,f}$ such that $\nu(C^1_K \times_{\hat{\mathcal{O}}^*} \hat{\mathcal{O}}) = 1$ and $\nu(\mathfrak{a}\,\cdot) = N(\mathfrak{a})^{-\beta}\nu$ for all $\mathfrak{a} \in I$; namely, the measure corresponding to a point $x \in C^1_K \times_{\hat{\mathcal{O}}^*} \hat{\mathcal{O}}^*$ is given by $\zeta_K(\beta)^{-1} \sum_{\mathfrak{a} \in I} N(\mathfrak{a})^{-\beta}\delta_{\mathfrak{a}x}$.*

5.5. In the global function field case with field of constants $\mathbb{F}_q$ the equidistribution result analogous to Theorem 3.1 fails because the L-function attached to the trivial Grössencharacter has an infinite set of simple poles on the line $\mathrm{Re}(s) = 1$ at $s = 1 + \frac{2\pi ik}{\log q}$. These extra poles produce oscillations in the counting function of the Dirichlet series coefficients below x. For geometric extensions the pole locations and residues for the trivial character do not change on the line $\mathrm{Re}(s) = 1$, and the ergodicity result can still be obtained in this case. However for non-geometric extensions, the change to the ground field $\mathbb{F}_{q^m}$ produces an L-function for the trivial character having a closer spacing of poles on the line $\mathrm{Re}(s) = 1$.

Assume K is a global function field and L is a finite abelian extension of K unramified at all finite places $\mathfrak{p} \notin S(\mathfrak{m})$. As in Theorem 4.2 consider the action of $I_{\mathfrak{m}}$ on $\mathbb{Z} \times \mathrm{Gal}(L/K) \times \mathbb{A}_{S(\mathfrak{m})}/\hat{\mathcal{O}}^*_{S(\mathfrak{m})}$, given by

$$\mathfrak{a}(n, g, x) = (n + \deg \mathfrak{a}, \sigma(\mathfrak{a})g, \mathfrak{a}x).$$

If the extension L/K is not geometric, then this action is *not* ergodic, since the image Γ of $I_{\mathfrak{m}}$ in $\mathbb{Z} \times \mathrm{Gal}(L/K)$ under the homomorphism $\mathfrak{a} \mapsto (\deg \mathfrak{a}, \sigma(\mathfrak{a}))$ is a proper subgroup. Namely, if the constant field of L is $\mathbb{F}_{q^m}$, then Γ is the kernel of the homomorphism

$$\mathbb{Z} \times \mathrm{Gal}(L/K) \to \mathrm{Gal}(\mathbb{F}_{q^m}/\mathbb{F}_q), \ \ (n, g) \mapsto (g|_{\mathbb{F}_{q^m}})\,\mathrm{Frob}^{-n}_{\mathbb{F}_{q^m}/\mathbb{F}_q}\,.$$

Nevertheless Theorem 4.2 can be formulated by saying that the kernel of the homomorphism $I_\mathfrak{m} \to \mathbb{Z} \times \mathrm{Gal}(L/K)$, $\mathfrak{a} \mapsto (\deg \mathfrak{a}, \sigma(\mathfrak{a}))$, acts ergodically on $\mathbb{A}_{S(\mathfrak{m})}/\hat{\mathcal{O}}^*_{S(\mathfrak{m})}$, and in this form the theorem remains true for non-geometric extensions. To put it differently, the asymptotic range of the cocycle c defined in the proof of Theorem 4.2 is exactly the subgroup $\Gamma \subset \mathbb{Z} \times \mathrm{Gal}(L/K)$. This is basically a reformulation of results of [17, Section 2], and this can also be proved using the same arguments as in our proof of Theorem 4.2.

Acknowledgments. This work came about through the authors' participation at the conference *Noncommutative Geometry: Multiple Connections*, held at Ohio State University, May 2012. The conference was supported by the Mathematics Research Institute, the NSF, and Ohio State University.

REFERENCES

[1] H. Araki and E.J. Woods, *A classification of factors*, Publ. Res. Inst. Math. Sci. Ser. A **4** (1968/1969), 51–130.
[2] B. Blackadar, *The regular representation of restricted direct product groups*, J. Func. Anal. **25** (1977), no. 3, 267–274.
[3] J.-B. Bost and A. Connes, *Hecke algebras, type III factors and phase transitions with spontaneous symmetry breaking in number theory*, Selecta Math. (N.S.) **1** (1995), no. 3, 411–457.
[4] A. Connes, *Trace formula in noncommutative geometry and the zeros of the Riemann zeta function*, Selecta Math. (N.S.) **5** (1999), no. 1, 29–106.
[5] J. Feldman and C.C. Moore, *Ergodic equivalence relations, cohomology, and von Neumann algebras. I*, Trans. Amer. Math. Soc. **234** (1977), no. 2, 289–324.
[6] M.D. Fried and M. Jarden, *Field arithmetic*, A Series of Modern Surveys in Mathematics, **11**, Springer-Verlag, Berlin, 2008.
[7] E. Hecke, *Eine neue Art von Zetafunktionen und ihre Beziehungen zur Verteilung der Primzahlen (Erste Meitteilung)*, Math. Zeitschrift **1** (1918), 357–376.
[8] E. Hecke, *Eine neue Art von Zetafunktionen und ihre Beziehungen zur Verteilung der Primzahlen (Zweite Meitteilung)*, Math. Zeitschrift **6** (1920), 11–51.
[9] W. Krieger, *On the Araki-Woods asymptotic ratio set and non-singular transformations of a measure space*, in "Contributions to Ergodic Theory and Probability" (Proc. Conf., Ohio State Univ., Columbus, Ohio, 1970), 158–177, Lecture Notes in Math., Vol. **160**, Springer, Berlin, 1970.
[10] M. Laca, N.S. Larsen and S. Neshveyev, *On Bost-Connes type systems for number fields*, J. Number Theory **129** (2009), no. 2, 325–338.
[11] M. Laca and I. Raeburn, *The ideal structure of the Hecke C^*-algebra of Bost and Connes*, Math. Ann. **318** (2000), no. 3, 433–451.
[12] S. Lang, *Algebraic Number Theory. Second Edition*, Springer, New York 1990. (First Edition: Addison-Wesley 1970).
[13] T. Mitsui, *Generalized prime number theorem*, Jap. J. Math. **26** (1956), 1–42.
[14] W. Narkiewicz, *Elementary and analytic theory of algebraic numbers. Third edition*, Springer Monographs in Mathematics, Springer-Verlag, Berlin, 2004. (Second Edition, 1990).
[15] S. Neshveyev, *Ergodicity of the action of the positive rationals on the group of finite adeles and the Bost-Connes phase transition theorem*, Proc. Amer. Math. Soc. **130** (2002), no. 10, 2999–3003.
[16] S. Neshveyev, *Von Neumann algebras arising from Bost-Connes type systems*, Int. Math. Res. Not. IMRN **2011**, no. 1, 217–236.
[17] S. Neshveyev and S. Rustad, *Bost-Connes systems associated with function fields*, preprint arXiv:1112.5826v2 [math.OA].
[18] J. Neukirch, *Algebraic number theory*, Fundamental Principles of Mathematical Sciences, **322**, Springer-Verlag, Berlin, 1999.
[19] J. Tate, *Fourier Analysis in Number Fields and Hecke's Zeta Functions*, Thesis: Princeton, May 1950. Reprinted in J. W. S. Cassels and A. Fröhlich, Eds., *Algebraic Number Theory*, 305–347, Academic Press, London, 1967.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MICHIGAN, ANN ARBOR, MI 48109-1043, USA
E-mail address: `lagarias@umich.edu`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF OSLO, P.O. BOX 1053 BLINDERN, N-0316 OSLO, NORWAY
E-mail address: `sergeyn@math.uio.no`